



# Active Bacterial Core Surveillance (ABCs) Report

## Emerging Infections Program Network

### Methicillin-Resistant *Staphylococcus aureus*, 2013



#### ABCs Areas

California (3 county San Francisco Bay area); Colorado (5 county Denver area); Connecticut; Georgia (8 county Atlanta area); Maryland (Baltimore City and County); Minnesota (2 metro Twin City counties); New York (1 Rochester county); Oregon (3 county Portland area); Tennessee (1 Nashville county).

#### ABCs Population

The surveillance areas represent **19,842,441** persons

Source: National Center for Health Statistics bridged-race vintage 2013 postcensal file.

#### ABCs Case Definition

Invasive methicillin-resistant *Staphylococcus aureus* (MRSA) disease: isolation of MRSA from a normally sterile site in a resident of the surveillance area in 2013. Cases of disease are classified into one of three epidemiologic classifications. A case is classified as hospital-onset (HO) if the MRSA culture was obtained on or after the fourth calendar day of hospitalization, where admission is hospital day 1; as healthcare-associated community-onset (HACO) if the culture was obtained in an outpatient setting or before the fourth calendar day of hospitalization and had one or more of the following: 1) a history of hospitalization, surgery, dialysis, or residence in a long term care facility in the previous year, or 2) the presence of a central vascular catheter (CVC) within 2 days prior to MRSA culture; and as community-associated (CA) if none of the previously mentioned criteria are met.

#### ABCs Methodology

ABCs personnel routinely contacted all microbiology laboratories serving healthcare facilities in their area to identify cases. Standardized case report forms that include information on demographic characteristics, clinical syndrome, and outcome of illness were completed for each identified case. Convenience samples of isolates were collected and sent to CDC for routine testing, including antimicrobial susceptibility testing, toxin testing, *SCCmec* typing, and *spa* typing. Pulsed field gel electrophoresis (PFGE) of all isolates was discontinued in 2008; up until 2012, PFGE was inferred based on a validated algorithm (<http://www.cdc.gov/HAI/settings/lab/inferred-PFGE-algorithm.html>). Starting in 2012, *spa* typing was added to the routine laboratory testing. Pulsed field type is currently inferred based on *spa* type, inferred MLST clonal complex and molecular characteristics of the isolates (<http://www.cdc.gov/HAI/settings/lab/CCalgorithm.html>). In 2013, isolates were only collected in five sites (California, Georgia, Minnesota, New York, and Tennessee). Regular laboratory audits were performed to ensure completeness of case detection.

Rates of invasive MRSA disease among all patients were calculated using population estimates for 2013. Cases with unknown race were assigned race based on distribution of known age, race, and gender by EIP site. Confidence intervals for nationally estimated incidence rates of disease and mortality were calculated based on the gamma distribution (Stat Med, 1997 16:791-801).

Rates of invasive MRSA disease among patients who were undergoing chronic dialysis treatment were calculated using the December 31, 2012 point prevalent counts of patients on dialysis from the United States Renal Data System (USRDS) (<http://www.usrds.org/adr.htm>).

#### ABCs Results

##### Reported Race among 4576 Cases

Race	No. (Rate <sup>a</sup> )
White	2843 (20.6)
Black	1563 (40.6)
Other	170 (7.9)

Unknown race (n=390) distributed amongst known

<sup>a</sup> Cases per 100,000 population for ABCs areas (crude rates)

##### Reported Cases on Chronic Dialysis (n=872)

Dialysis and Access Type	No. (%)
Type of dialysis	
Peritoneal	29 (3.3)
Hemodialysis	834 (95.6)
AV Fistula/Graft	340 (39.0)
CVC	462 (53.0)
Unknown	32 (3.7)
Unknown	9 (1.0)

##### Cases, Deaths & Inferred PFGE type by Epidemiological Classification

MRSA Class	No. (Rate) Cases <sup>b</sup>	No. (Rate) Deaths <sup>c</sup>	Inferred PFGE Type (%)			
			Tot N	USA100	USA300	USA500/Iberian
CA	998 (5.0)	102 (0.5)	150	22 (14.7)	103 (68.7)	6 (4.0)
HCA <sup>a</sup>	3529 (17.8)	480 (2.4)	554	243 (43.9)	165 (29.8)	75 (13.5)
HO	704 (3.6)	135 (0.7)	111	48 (43.2)	41 (36.9)	9 (8.1)
HACO	2825 (14.2)	345 (1.7)	443	195 (44.0)	124 (28.0)	66 (14.9)

<sup>a</sup> HCA: Healthcare-associated invasive MRSA infection; sum of patients that are classified as either the HO or HACO classes

<sup>b</sup> n= 49 epidemiologic category unknown

<sup>c</sup> n=6; epidemiologic category unknown

## Reported Clinical Syndrome by Epidemiological Class

Syndrome <sup>a</sup>	CA (n=998)	HACO (n=2825)	HO (n=704)
	No. (%)	No. (%)	No. (%)
Bloodstream Infection <sup>b</sup>			
with other syndrome	521 (52.2)	1363 (48.2)	216 (30.7)
with no other syndrome	210 (21.0)	963 (34.1)	247 (35.1)
Pneumonia	141 (14.1)	349 (12.4)	95 (13.5)
Lower Respiratory Infection <sup>c</sup>	44 (4.4)	103 (3.6)	28 (4.0)
Osteomyelitis	154 (15.4)	418 (14.8)	142 (20.2)
Endocarditis	75 (7.5)	168 (5.9)	29 (4.1)
Cellulitis	180 (18.0)	233 (8.2)	56 (8.0)
Wounds			
Surgical <sup>d</sup>	8 (0.8)	136 (4.8)	28 (4.0)
Decubitus/Pressure Ulcers	10 (1.0)	53 (1.9)	11 (1.6)
Skin Abscesses <sup>e</sup>	94 (9.4)	87 (3.1)	18 (2.6)
Other Wounds <sup>f</sup>	32 (3.2)	96 (3.4)	32 (4.5)
Traumatic	11 (1.1)	6 (0.2)	3 (0.4)

<sup>a</sup> Some case patients had more than one syndrome.

<sup>b</sup> Catheter site infection or AV fistula infection only are included in BSI with other syndrome.


<sup>c</sup> Lower Respiratory Infection is defined as: a patient with pneumonia documented in their discharge summary who has a positive MRSA non-sterile respiratory specimen with accompanying chest radiology results documenting any of the following: bronchopneumonia/pneumonia, air space density/opacity, new or changed infiltrates.

<sup>d</sup> Combines deep tissue/organ infection and infection of a surgical wound, post operatively.

<sup>e</sup> Category includes skin abscess, necrotizing fasciitis, gangrene.

<sup>f</sup> Category includes non-traumatic and other chronic wound infections.

**Incidence of Invasive MRSA, by  
Epidemiological Class and Age Group  
2013**



## National Estimates and Adjusted Incidence Rates of Invasive MRSA Infections

Epidemiologic Category	Estimated Cases of Infection					
	Non-Dialysis Patients		Dialysis Patients		Total	
	Estimated No.	Incidence Rate (Confidence Interval) <sup>a</sup>	Estimated No.	Incidence Rate (Confidence Interval) <sup>b</sup>	Estimated No.	Incidence Rate (Confidence Interval) <sup>d</sup>
CA	16,117	5.10 (3.92-6.74)	NA	NA	16,117	5.10 (3.92-6.74)
HCA	43,864	13.90 (12.02-16.23)	13,919	3079.89 (2331.21-4076.80)	57,783	18.28 (15.34-22.04)
HO	9,820	3.11 (2.24-4.39)	1,584	350.44 (153.54-710.62)	11,404	3.61 (2.46-5.40)
HACO	34,044	10.78 (9.21-12.76)	12,335	2729.00 (2026.78-3681.88)	46,379	14.67 (12.10-18.00)
Overall <sup>c</sup>	60,772	19.25 (16.94-22.05)	13,921	3079.89 (2331.21-4076.80)	74,693	23.63 (20.24-27.84)

<sup>a</sup> National Estimates and Incidence (no. per 100,000 population per year) are adjusted for age, race, gender and receipt of chronic dialysis using 2013 US Census Data.

<sup>b</sup> National Estimates and Incidence (no. per 100,000 dialysis patients per year) for dialysis patients are adjusted for age, race and gender using 2012 USRDS point prevalence data.

<sup>c</sup> 49 cases could not be classified into an epidemiological category or category is unknown and therefore are counted in the overall estimate only.

<sup>d</sup> Starting in 2011, confidence intervals on national estimates were determined for each of the 72 age/race/gender/dialysis specific strata and summarized for an overall national estimate, accounting for variance across all strata producing a more conservative estimate (with wider confidence intervals) compared to estimates prior to 2011.

### National Estimates and Adjusted Incidence Rates for Mortality among Cases

Epidemiologic Class	Estimated No.	Mortality Rate
		(Confidence Interval) <sup>a</sup>
CA	1,710	0.54 (0.28-1.00)
HCA	8,150	2.58 (1.68-4.01)
HO	2,331	0.74 (0.33-1.59)
HACO	5,819	1.84 (1.12-3.02)
Overall <sup>b</sup>	9,937	3.14 (2.14-4.68)

<sup>a</sup> National Estimates and Mortality Rate (no. per 100,000 population per year) are adjusted for age, race, gender and receipt of chronic dialysis using 2013 US Census Data

<sup>b</sup> 49 cases could not be classified into an epidemiological category or category is unknown and therefore are counted in the overall estimate **only**.

### National Metric for Healthy People 2020 and the Department of Health and Human Services Action Plan to Prevent Healthcare-Associated Infections

	Disease Rate			Estimate of Cases in United States. <sup>a</sup>		
	Baseline (07-08)	2013	% Change	Baseline (07-08)	2013	Difference
HCA	27.08	18.28	-32.50	82,000	58,000	24,000

<sup>a</sup> Disease Rate (no. per 100,000 population per year) and National Estimates are adjusted for age, race, gender and receipt of chronic dialysis using 2012 US Census Data

### ABCs Discussion

Surveillance data from 2013 represent the ninth full year of performing population-based surveillance for invasive MRSA infections through the Emerging Infections Program/Active Bacterial Core Surveillance Activity.

Compared to 2012, incidence of healthcare-associated MRSA decreased by 1.01% while incidence of community-associated MRSA increased by 6.51%. Compared to the baseline incidence (2007-2008 calendar years) identified in the HHS Action Plan, there was a decrease of 32.50% for healthcare-associated MRSA.

### Citation

1. Centers for Disease Control and Prevention. 2013. Active Bacterial Core Surveillance Report, Emerging Infections Program Network, Methicillin-Resistant *Staphylococcus aureus*, 2013.

Available via the Internet: <http://www.cdc.gov/abcs/reports-findings/survreports/mrsa13.html>

For more information, visit our web sites: <http://www.cdc.gov/abcs/index.html>, <http://www.cdc.gov/mrsa>